


**G.GPE.B.4: Quadrilaterals in the Coordinate Plane 1**


- 1 On the set of axes below, the coordinates of three vertices of trapezoid  $ABCD$  are  $A(2,1)$ ,  $B(5,4)$ , and  $D(-2,3)$ .




Which point could be vertex  $C$ ?

- 1)  $(1,5)$
  - 2)  $(4,10)$
  - 3)  $(-1,6)$
  - 4)  $(-3,8)$
- 2 A quadrilateral has vertices with coordinates  $(-3,1)$ ,  $(0,3)$ ,  $(5,2)$ , and  $(-1,-2)$ . Which type of quadrilateral is this?
- 1) rhombus
  - 2) rectangle
  - 3) square
  - 4) trapezoid
- 3 The coordinates of the vertices of parallelogram  $CDEH$  are  $C(-5,5)$ ,  $D(2,5)$ ,  $E(-1,-1)$ , and  $H(-8,-1)$ . What are the coordinates of  $P$ , the point of intersection of diagonals  $\overline{CE}$  and  $\overline{DH}$ ?
- 1)  $(-2,3)$
  - 2)  $(-2,2)$
  - 3)  $(-3,2)$
  - 4)  $(-3,-2)$
- 4 Rectangle  $ABCD$  has two vertices at coordinates  $A(-1,-3)$  and  $B(6,5)$ . The slope of  $\overline{BC}$  is
- 1)  $-\frac{7}{8}$
  - 2)  $\frac{7}{8}$
  - 3)  $-\frac{8}{7}$
  - 4)  $\frac{8}{7}$
- 5 Parallelogram  $ABCD$  has coordinates  $A(0,7)$  and  $C(2,1)$ . Which statement would prove that  $ABCD$  is a rhombus?
- 1) The midpoint of  $\overline{AC}$  is  $(1,4)$ .
  - 2) The length of  $\overline{BD}$  is  $\sqrt{40}$ .
  - 3) The slope of  $\overline{BD}$  is  $\frac{1}{3}$ .
  - 4) The slope of  $\overline{AB}$  is  $\frac{1}{3}$ .
- 6 The diagonals of rhombus  $TEAM$  intersect at  $P(2,1)$ . If the equation of the line that contains diagonal  $\overline{TA}$  is  $y = -x + 3$ , what is the equation of a line that contains diagonal  $\overline{EM}$ ?
- 1)  $y = x - 1$
  - 2)  $y = x - 3$
  - 3)  $y = -x - 1$
  - 4)  $y = -x - 3$


- 7 In square  $GEOM$ , the coordinates of  $G$  are  $(2, -2)$  and the coordinates of  $O$  are  $(-4, 2)$ . Determine and state the coordinates of vertices  $E$  and  $M$ . [The use of the set of axes below is optional.]




- 9 Quadrilateral  $NATS$  has coordinates  $N(-4, -3)$ ,  $A(1, 2)$ ,  $T(8, 1)$ , and  $S(3, -4)$ . Prove quadrilateral  $NATS$  is a rhombus. [The use of the set of axes below is optional.]




- 8 The coordinates of the vertices of quadrilateral  $HYPE$  are  $H(-3, 6)$ ,  $Y(2, 9)$ ,  $P(8, -1)$ , and  $E(3, -4)$ . Prove  $HYPE$  is a rectangle. [The use of the set of axes below is optional.]




- 10 Parallelogram  $MATH$  has vertices  $M(-7,-2)$ ,  $A(0,4)$ ,  $T(9,2)$ , and  $H(2,-4)$ . Prove that parallelogram  $MATH$  is a rhombus. [The use of the set of axes below is optional.] Determine and state the area of  $MATH$ .




- 11 Quadrilateral  $PQRS$  has vertices  $P(-2,3)$ ,  $Q(3,8)$ ,  $R(4,1)$ , and  $S(-1,-4)$ . Prove that  $PQRS$  is a rhombus. Prove that  $PQRS$  is *not* a square. [The use of the set of axes below is optional.]




- 12 The coordinates of the vertices of quadrilateral  $ABCD$  are  $A(0,4)$ ,  $B(3,8)$ ,  $C(8,3)$ , and  $D(5,-1)$ . Prove that  $ABCD$  is a parallelogram, but *not* a rectangle. [The use of the set of axes below is optional.]




- 13 The vertices of quadrilateral  $MATH$  have coordinates  $M(-4,2)$ ,  $A(-1,-3)$ ,  $T(9,3)$ , and  $H(6,8)$ . Prove that quadrilateral  $MATH$  is a parallelogram. Prove that quadrilateral  $MATH$  is a rectangle. [The use of the set of axes below is optional.]




- 14 Riley plotted  $A(-1,6)$ ,  $B(3,8)$ ,  $C(6,-1)$ , and  $D(1,0)$  to form a quadrilateral. Prove that Riley's quadrilateral  $ABCD$  is a trapezoid. [The use of the set of axes below is optional.] Riley defines an isosceles trapezoid as a trapezoid with congruent diagonals. Use Riley's definition to prove that  $ABCD$  is *not* an isosceles trapezoid.




- 15 Quadrilateral  $MATH$  has vertices with coordinates  $M(-1,7)$ ,  $A(3,5)$ ,  $T(2,-7)$ , and  $H(-6,-3)$ . Prove that quadrilateral  $MATH$  is a trapezoid. State the coordinates of point Y such that point  $A$  is the midpoint of  $\overline{MY}$ . Prove that quadrilateral  $MYTH$  is a rectangle. [The use of the set of axes below is optional.]




- 16 In the coordinate plane, the vertices of  $\triangle RST$  are  $R(6, -1)$ ,  $S(1, -4)$ , and  $T(-5, 6)$ . Prove that  $\triangle RST$  is a right triangle. State the coordinates of point  $P$  such that quadrilateral  $RSTP$  is a rectangle. Prove that your quadrilateral  $RSTP$  is a rectangle. [The use of the set of axes below is optional.]




- 17 In the coordinate plane, the vertices of triangle  $PAT$  are  $P(-1, -6)$ ,  $A(-4, 5)$ , and  $T(5, -2)$ . Prove that  $\triangle PAT$  is an isosceles triangle. State the coordinates of  $R$  so that quadrilateral  $PART$  is a parallelogram. Prove that quadrilateral  $PART$  is a parallelogram. [The use of the set of axes below is optional.]



- 18 The coordinates of the vertices of  $\triangle ABC$  are  $A(1,2)$ ,  $B(-5,3)$ , and  $C(-6,-3)$ . Prove that  $\triangle ABC$  is isosceles. State the coordinates of point  $D$  such that quadrilateral  $ABCD$  is a square. Prove that your quadrilateral  $ABCD$  is a square. [The use of the set of axes below is optional.]



- 19 The coordinates of the vertices of  $\triangle ABC$  are  $A(-2,4)$ ,  $B(-7,-1)$ , and  $C(-3,-3)$ . Prove that  $\triangle ABC$  is isosceles. State the coordinates of  $\triangle A'B'C'$ , the image of  $\triangle ABC$ , after a translation 5 units to the right and 5 units down. Prove that quadrilateral  $AA'C'C$  is a rhombus. [The use of the set of axes below is optional.]




- 20 Given: Triangle  $DUC$  with coordinates  $D(-3, -1)$ ,  $U(-1, 8)$ , and  $C(8, 6)$

Prove:  $\triangle DUC$  is a right triangle


Point  $U$  is reflected over  $\overline{DC}$  to locate its image point,  $U'$ , forming quadrilateral  $DUCU'$ .

Prove quadrilateral  $DUCU'$  is a square.

[The use of the set of axes below is optional.]



- 21 In rhombus  $MATH$ , the coordinates of the endpoints of the diagonal  $\overline{MT}$  are  $M(0, -1)$  and  $T(4, 6)$ . Write an equation of the line that contains diagonal  $\overline{AH}$ . [Use of the set of axes below is optional.] Using the given information, explain how you know that your line contains diagonal  $\overline{AH}$ .





**G.GPE.B.4: Quadrilaterals in the Coordinate Plane 1**  
**Answer Section**

1 ANS: 4

$$m_{\overline{AD}} = \frac{3-1}{-2-2} = \frac{2}{-4} = -\frac{1}{2} \quad \text{A pair of opposite sides is parallel.}$$

$$m_{\overline{BC}} = \frac{8-4}{-3-5} = \frac{4}{-8} = -\frac{1}{2}$$

REF: 082321geo

2 ANS: 4

$$\frac{-2-1}{-1--3} = \frac{-3}{2} \quad \frac{3-2}{0-5} = \frac{1}{-5} \quad \frac{3-1}{0--3} = \frac{2}{3} \quad \frac{2--2}{5--1} = \frac{4}{6} = \frac{2}{3}$$

REF: 081522geo

3 ANS: 3

$$M_x = \frac{-5+-1}{2} = -\frac{6}{2} = -3 \quad M_y = \frac{5+-1}{2} = \frac{4}{2} = 2$$

REF: 081902geo

4 ANS: 1

$$m_{\overline{AB}} = \frac{-3-5}{-1-6} = \frac{-8}{-7} = \frac{8}{7}$$

REF: 062315geo

5 ANS: 3

$$\frac{7-1}{0-2} = \frac{6}{-2} = -3 \quad \text{The diagonals of a rhombus are perpendicular.}$$

REF: 011719geo

6 ANS: 1


$$m_{\overline{TA}} = -1 \quad y = mx + b$$

$$m_{\overline{EM}} = 1 \quad 1 = 1(2) + b$$

$$-1 = b$$


REF: 081614geo

7 ANS:



REF: 011731geo


8 ANS:



1) Quadrilateral *HYPE* with  $H(-3,6)$ ,  $Y(2,9)$ ,  $P(8,-1)$ , and  $E(3,-4)$  (Given); 2) Slope of  $\overline{HY}$  and  $\overline{PE}$  is  $\frac{3}{5}$ , slope of  $\overline{YP}$  and  $\overline{EH}$  is  $-\frac{5}{3}$  (Slope determined graphically); 3)  $\overline{HY} \perp \overline{YP}$ ,  $\overline{PE} \perp \overline{EH}$ ,  $\overline{YP} \perp \overline{PE}$ ,  $\overline{EH} \perp \overline{HY}$  (The slopes of perpendicular lines are opposite reciprocals); 4)  $\angle H$ ,  $\angle Y$ ,  $\angle P$ ,  $\angle E$  are right angles (Perpendicular lines form right angles); 5) *HYPE* is a rectangle (A rectangle has four right angles).

REF: 082233geo

9 ANS:



$$\overline{AN} \cong \overline{AT} \cong \overline{TS} \cong \overline{SN}$$

Quadrilateral *NATS* is a rhombus

$$\sqrt{5^2 + 5^2} = \sqrt{7^2 + 1^2} = \sqrt{5^2 + 5^2} = \sqrt{7^2 + 1^2}$$


$$\sqrt{50} = \sqrt{50} = \sqrt{50} = \sqrt{50}$$

because all four sides are congruent.

REF: 012032geo

10 ANS:

A rhombus has four congruent sides. Since each side measures  $\sqrt{85}$ , all four sides of *MATH* are congruent, and




*MATH* is a rhombus.  $16 \times 8 - (21 + 9 + 21 + 9) = 68$

REF: 062334geo

11 ANS:


$\overline{PQ} \sqrt{(8-3)^2 + (3--2)^2} = \sqrt{50}$ 
 $\overline{QR} \sqrt{(1-8)^2 + (4-3)^2} = \sqrt{50}$ 
 $\overline{RS} \sqrt{(-4-1)^2 + (-1-4)^2} = \sqrt{50}$   
 $\overline{PS} \sqrt{(-4-3)^2 + (-1--2)^2} = \sqrt{50}$ 
*PQRS* is a rhombus because all sides are congruent.  $m_{\overline{PQ}} = \frac{8-3}{3--2} = \frac{5}{5} = 1$   
 $m_{\overline{QR}} = \frac{1-8}{4-3} = -7$  Because the slopes of adjacent sides are not opposite reciprocals, they are not perpendicular



and do not form a right angle. Therefore *PQRS* is not a square.

REF: 061735geo


12 ANS:



$\overline{AD}$  and  $\overline{BC}$  have equal slope, so are parallel.  $\overline{AB}$  and  $\overline{CD}$  have equal slope, so are parallel. Since both pairs of opposite sides are parallel, *ABCD* is a parallelogram. The slope of  $\overline{AB}$  and  $\overline{BC}$  are not opposite reciprocals, so they are not perpendicular, and so  $\angle B$  is not a right angle. *ABCD* is not a rectangle since all four angles are not right angles.

REF: 082334geo

13 ANS:




$$m_{\overline{MH}} = \frac{6}{10} = \frac{3}{5}, m_{\overline{AT}} = \frac{6}{10} = \frac{3}{5}, m_{\overline{MA}} = -\frac{5}{3}, m_{\overline{HT}} = -\frac{5}{3}; \overline{MH} \parallel \overline{AT} \text{ and } \overline{MA} \parallel \overline{HT}.$$

$MATH$  is a parallelogram since both sides of opposite sides are parallel.  $m_{\overline{MA}} = -\frac{5}{3}, m_{\overline{AT}} = \frac{3}{5}$ . Since the slopes are negative reciprocals,  $\overline{MA} \perp \overline{AT}$  and  $\angle A$  is a right angle.  $MATH$  is a rectangle because it is a parallelogram with a right angle.

REF: 081835geo

14 ANS:



$$m_{\overline{AD}} = \frac{0-6}{1- -1} = -3 \quad \overline{AD} \parallel \overline{BC} \text{ because their slopes are equal. } ABCD \text{ is a trapezoid}$$

$$m_{\overline{BC}} = \frac{-1-8}{6-3} = -3$$


because it has a pair of parallel sides.  $AC = \sqrt{(-1-6)^2 + (6- -1)^2} = \sqrt{98}$   $ABCD$  is not an isosceles trapezoid

$$BD = \sqrt{(8-0)^2 + (3-1)^2} = \sqrt{68}$$

because its diagonals are not congruent.

REF: 061932geo

15 ANS:



The slope of  $\overline{MA}$  and  $\overline{TH}$  equals  $-\frac{1}{2}$ . Distinct lines with equal slope are parallel.  $MATH$  is a trapezoid because it has a pair of parallel lines.  $(7,3)$ . The slope of  $\overline{MY}$  and  $\overline{TH}$  equals  $-\frac{1}{2}$ . The slope of  $\overline{YT}$  and  $\overline{HM}$  equals 2. The slopes of each side are opposite reciprocals and therefore perpendicular. Perpendicular sides form right angles, so  $MYTH$  has four right angles and is a rectangle.


REF: 012435geo

16 ANS:

$m_{\overline{TS}} = \frac{-10}{6} = -\frac{5}{3}$   $m_{\overline{SR}} = \frac{3}{5}$  Since the slopes of  $\overline{TS}$  and  $\overline{SR}$  are opposite reciprocals, they are perpendicular and

form a right angle.  $\triangle RST$  is a right triangle because  $\angle S$  is a right angle.  $P(0,9)$   $m_{\overline{RP}} = \frac{-10}{6} = -\frac{5}{3}$   $m_{\overline{PT}} = \frac{3}{5}$

Since the slopes of all four adjacent sides ( $\overline{TS}$  and  $\overline{SR}$ ,  $\overline{SR}$  and  $\overline{RP}$ ,  $\overline{PT}$  and  $\overline{TS}$ ,  $\overline{RP}$  and  $\overline{PT}$ ) are opposite reciprocals, they are perpendicular and form right angles. Quadrilateral  $RSTP$  is a rectangle because it has four right angles.




REF: 061536geo

17 ANS:

$\triangle PAT$  is an isosceles triangle because sides  $\overline{AP}$  and  $\overline{AT}$  are congruent ( $\sqrt{3^2 + 11^2} = \sqrt{7^2 + 9^2} = \sqrt{130}$ ).

$R(2,9)$ . Quadrilateral  $PART$  is a parallelogram because the opposite sides are parallel since they have equal slopes



$$(m_{\overline{AR}} = \frac{4}{6} = \frac{2}{3}; m_{\overline{PT}} = \frac{4}{6} = \frac{2}{3}; m_{\overline{PA}} = -\frac{11}{3}; m_{\overline{RT}} = -\frac{11}{3})$$


REF: 011835geo

18 ANS:

$AB = \sqrt{(-5-1)^2 + (3-2)^2} = \sqrt{37}$ ,  $BC = \sqrt{(-5-(-6))^2 + (3-(-3))^2} = \sqrt{37}$  (because  $AB = BC$ ,  $\triangle ABC$  is

isosceles).  $(0, -4)$ .  $AD = \sqrt{(1-0)^2 + (2-(-4))^2} = \sqrt{37}$ ,  $CD = \sqrt{(-6-0)^2 + (-3-(-4))^2} = \sqrt{37}$ ,

$m_{\overline{AB}} = \frac{3-2}{-5-1} = -\frac{1}{6}$ ,  $m_{\overline{CB}} = \frac{3-(-3)}{-5-(-6)} = 6$  ( $ABCD$  is a square because all four sides are congruent, consecutive sides



are perpendicular since slopes are opposite reciprocals and so  $\angle B$  is a right angle).

REF: 081935geo

19 ANS:


$$\sqrt{(-2 - -7)^2 + (4 - -1)^2} = \sqrt{(-2 - -3)^2 + (4 - -3)^2} \text{ Since } \overline{AB} \text{ and } \overline{AC} \text{ are congruent, } \triangle ABC \text{ is isosceles.}$$

$$\sqrt{50} = \sqrt{50}$$

$$A'(3, -1), B'(-2, -6), C'(2, -8). \quad AC = \sqrt{50} \quad AA' = \sqrt{(-2 - 3)^2 + (4 - -1)^2}, \quad A'C' = \sqrt{50} \text{ (translation preserves distance),}$$

$$CC' = \sqrt{(-3 - 2)^2 + (-3 - -8)^2} \text{ Since all four sides are congruent, } AA'C'C \text{ is a rhombus.}$$


$$= \sqrt{50}$$



REF: 062235geo

20 ANS:

$m_{\overline{DU}} = \frac{9}{2}$   $m_{\overline{UC}} = -\frac{2}{9}$  Since the slopes of  $\overline{DU}$  and  $\overline{UC}$  are opposite reciprocals, they are perpendicular and form a right angle.  $\triangle DUC$  is a right triangle because  $\angle DUC$  is a right angle. Each side of quadrilateral  $DUCU'$  is  $\sqrt{9^2 + 2^2} = \sqrt{85}$ . Quadrilateral  $DUCU'$  is a square because all four sides are congruent and it has a right angle.



REF: 012335geo

21 ANS:

$M\left(\frac{4+0}{2}, \frac{6-1}{2}\right) = M\left(2, \frac{5}{2}\right)$   $m = \frac{6 - -1}{4 - 0} = \frac{7}{4}$   $m_{\perp} = -\frac{4}{7}$   $y - 2.5 = -\frac{4}{7}(x - 2)$  The diagonals,  $\overline{MT}$  and  $\overline{AH}$ , of rhombus  $MATH$  are perpendicular bisectors of each other.

REF: fall1411geo